



DERMAL FILLER INSTRUCTIONS

Pre-Treatment Instructions:

- For optimal results, and to minimize the chance of bleeding or bruising at the injection site, please avoid all blood-thinning medications and supplements for one week prior to your appointment. This includes over-the-counter medication such as aspirin, Motrin, ibuprofen, and Aleve. Also avoid vitamin E, and omega-3 capsules. If you have a cardiovascular history, please check with your doctor prior to stopping use of aspirin.
- Inform your provider if you have a history of Cold Sores or Fever Blisters to receive advice on antiviral therapy prior to treatment. Valtrex 500 mg BID, 5 days, start 2 days prior to treatment.
- Begin taking Arnica (an herbal anti-inflammatory typically found in health food stores) to help the bruising and swelling at least 2 days prior to your appointment.
- Avoid topical products such as Tretinoin (Retin-A) retinols, retinoids, glycolic acid, alpha hydroxy acid, or any "anti-aging" products for two days before and after treatment.
- Do not drink alcoholic beverages 24 hours before or after your treatment to avoid extra bruising.
- Do not use dermal fillers if you are pregnant or breastfeeding

Post-Treatment Instructions:

- Avoid significant movement or massage of the treated area unless instructed by provider.
- Avoid strenuous exercise for 24 hours.
- Avoid extensive sun or heat for 72 hours.
- Wait a minimum of four weeks before skin care or laser treatments.
- Avoid consuming alcohol or salty food for 24-48 hours to avoid excessive swelling.
- If you have swelling, you may apply a cool compress for 15 minutes each hour.
- Use Tylenol (acetaminophen) for discomfort.
- Try to sleep face up and slightly elevated if you experience swelling.
- Apply Alastin Inhance four times daily to the injection sites.

Dermal fillers last, on average, 6-12 months. It is recommended at the time of your treatment to schedule your next appointment for optimal anti-aging benefits. The most typical side effects (bruising, swelling, redness) are temporary and will resolve. However, please call the office to notify us if you experience any additional side effects.